arXiv:1608.04877v1 [math.DG] 17 Aug 2016

ON KNOTTED SPHERES IN EUCLIDEAN 4-SPACE $\mathbb{E}^4$

Kadri Arslan

August 18, 2016

Abstract

In the present study we consider knotted spheres in Euclidean 4-space $\mathbb{E}^4$. Firstly, we give some basic curvature properties of knotted spheres in $\mathbb{E}^4$. Further, we obtained some results related with the conjugate nets and Laplace transforms of these kind of surfaces.

1 Introduction

[1]Let us consider a unit speed regular curve γ in $\mathbb{E}^4$ and a unit speed spherical curve ρ in $\mathbb{E}^2$. Then, the rotation of γ around ρ give rise a surface M in $\mathbb{E}^4$, which is called *rotational surface*. The rotational surfaces in $\mathbb{E}^4$ was first introduced by C. Moore in 1919 (see, [12]). Further, many researchers concentrated these studies on this subject, see for example [3], [4], [7] and [8]. The rotational surfaces in $\mathbb{E}^4$ with constant curvatures are studied in [13].

Let us denote the half-space $x_3 \geq 0, x_4 = 0$ by $\mathbb{E}^3_+(0)$ and take an arc α with the end point in the plane $x_3 = 0, x_4 = 0$ (denoted by Π). The rotation the half space plane $\mathbb{E}^3_+(0)$ by the angle v around the plane Π is denoted by $\mathbb{E}^3_+(v)$. Consequently, after the rotation the point with coordinates

[1]2010 AMS *Mathematics Subject Classification.* 53C40, 53C42

Key words and phrases: Rotational surface, Knotted sphere, Conjugate nets

x_1, x_2, x_3, x_4 passes into the point with the coordinates $\widetilde{x}_1, \widetilde{x}_2, \widetilde{x}_3, \widetilde{x}_4$ by

$$\begin{aligned} \widetilde{x}_1 &= x_1 \\ \widetilde{x}_2 &= x_2 \\ \widetilde{x}_3 &= x_3 \cos v - x_4 \sin v \\ \widetilde{x}_3 &= x_3 \sin v - x_4 \cos v. \end{aligned} \tag{1}$$

In rotation by 360° the points of α being in $\mathbb{E}^3_+(v)$, form the set M homeomorphic to S^2 [2]. Let α be a smooth curve with tangent vectors at p and q orthogonal to Π. Then M is a smooth surface which is called *knotted sphere* in $\mathbb{E}^4$ [1].

A net of curves on a surface M is called *conjugate*, if at every point the tangent directions of the curves of the net separate harmonically the asymptotic directions [11]. Consequently, for a surface M with a patch $X(u, v)$, a net of curves on M are conjugate if and only if the second partial derivative X_{uv} of X lies in the subspace spanned by X_u and X_v [10].

This paper is organized as follows: In section 2 we give some basic concepts of the second fundamental form and curvatures of the surfaces in $\mathbb{E}^4$. In Section 3 we consider knotted spheres in $\mathbb{E}^4$. Firstly, we give some basic curvature properties of knotted spheres in $\mathbb{E}^4$. Further, we introduce some kind of knotted spheres and obtained some results related with their curvatures. In section 4 we give some basic curvature properties of the conjugate nets on a surface in E^n. Further, we calculated the Laplace invariants and the Laplace transforms of the knotted sphere in $\mathbb{E}^4$.

2 Preliminaries

Let M be a local surface in $\mathbb{E}^n$ given with position vector $X(u, v)$. The tangent space T_pM is spanned by the vector fields X_u and X_v. In the chart (u, v) the coefficients of the first fundamental form of M are given by

$$E = \langle X_u, X_u \rangle, F = \langle X_u, X_v \rangle, G = \langle X_v, X_v \rangle, \tag{2}$$

where $\langle , \rangle$ is the inner product in $\mathbb{E}^n$. We assume that $X(u, v)$ is regular i.e., $W^2 = EG - F^2 \neq 0$ [9].

Consequently, the *Gaussian curvature* of M is given by

$$K = -\frac{1}{4W^2} \begin{vmatrix} E & E_u & E_v \\ F & F_u & F_v \\ G & G_u & G_v \end{vmatrix} - \frac{1}{2W} \left(\left(\frac{E_v - F_u}{W} \right)_v - \left(\frac{F_v - G_u}{W} \right)_u \right). \tag{3}$$

Let $\widetilde{\nabla}$ be the Riemannian connection of $\mathbb{E}^4$, and $X_1 = X_u$, $X_2 = X_v$ tangent vector fields of M then *Gauss equation* gives

$$\widetilde{\nabla}_{X_i} X_j = \sum_{k=1}^{2} \Gamma_{ij}^k X_k + h\left(X_i, X_j\right); \ 1 \leq i, j \leq 2, \tag{4}$$

where h is the second fundamental form and Γ_{ij}^k are the *Christoffel symbols* of M.

The *mean curvature vector* $\overrightarrow{H}$ of of M is given by

$$\overrightarrow{H} = \frac{1}{2W^2}\left(Eh(X_v, X_v) - 2Fh(X_u, X_v) + Gh(X_u, X_u)\right). \tag{5}$$

The norm of the mean curvature vector $\overrightarrow{H}$ is known as *mean curvature* of M. Recall that, a surface M is said to be *minimal* (resp. *flat*) if its mean curvature (resp. Gaussian curvature) vanishes identically [5].

3 Knotted Spheres in $\mathbb{E}^4$

Let M be a knotted sphere given with (1), then the position vector of M can be represented as:

$$X(u, v) = \left(x_1(u), x_2(u), x_3(u)\cos v - x_4(u)\sin v, x_3(u)\sin v + x_4(u)\cos v\right). \tag{6}$$

where

$$\gamma(u) = \left(x_1(u), x_2(u), x_3(u), x_4(u)\right),$$

is the profile curve of the surface [1]. Then, the tangent space T_pM of M is spanned by

$$\begin{aligned} X_u &= \left(x_1'(u), x_2'(u), x_3'(u)\cos v - x_4'(u)\sin v, x_3'(u)\sin v + x_4'(u)\cos v\right), \\ X_v &= \left(0, 0, -x_3(u)\sin v - x_4(u)\cos v, x_3(u)\cos v - x_4(u)\sin v\right). \end{aligned} \tag{7}$$

Consequently, the coefficients of first fundamental form become

$$\begin{aligned} E &= \langle X_u, X_u\rangle = 1, \\ F &= \langle X_u, X_v\rangle = x_3(u)x_4'(u) - x_3'(u)x_4(u), \\ G &= \langle X_v, X_v\rangle = x_3^2(u) + x_4^2(u). \end{aligned} \tag{8}$$

The Christoffel symbols Γ_{ij}^{k} of the canal surface M are given by

$$\begin{array}{lll} \Gamma_{11}^{1}=-\frac{FF_u}{W^2}, & \Gamma_{12}^{1}=-\frac{FG_u}{2W^2}, & \Gamma_{22}^{1}=-\frac{GG_u}{2W^2} \\ \Gamma_{11}^{2}=\frac{F_u}{W^2}, & \Gamma_{12}^{2}=\frac{G_u}{2W^2}, & \Gamma_{22}^{2}=\frac{FG_u}{2W^2}. \end{array} \tag{9}$$

which are symmetric with respect to the covariant indices ([9], p.398).

The second partial derivatives of r are expressed as follows:

$$\begin{aligned} X_{uu} &= (x_1''(u), x_2''(u), x_3''(u)\cos v - x_4''(u)\sin v, x_3''(u)\sin v + x_4''(u)\cos v)\,, \\ X_{uv} &= (0, 0, -x_3'(u)\sin v - x_4'(u)\cos v, x_3'(u)\cos v - x_4'(u)\sin v)\,, \\ X_{vv} &= (0, 0, -x_3(u)\cos v + x_4(u)\sin v, -x_3(u)\sin v - x_4(u)\cos v)\,. \end{aligned} \tag{10}$$

Using (7) with (10) we get

$$\begin{aligned} \langle X_{uu}, X_{vv}\rangle &= -\left(x_3(u)x_3''(u) + x_4(u)x_4''(u)\right), \\ \langle X_{vv}, X_u\rangle &= -\left(x_3(u)x_3'(u) + x_4(u)x_4'(u)\right), \\ \langle X_{uu}, X_v\rangle &= x_3(u)x_4''(u) - x_3''(u)x_4(u), \\ \langle r_{s\theta}, X_{uv}\rangle &= (x_3'(u))^2 + (x_4'(u))^2, \\ \langle X_{uu}, X_u\rangle &= 0, \\ \langle X_{vv}, X_v\rangle &= 0. \end{aligned} \tag{11}$$

Hence, taking into account (4), the Gauss equation implies the following equations;

$$\begin{aligned} \widetilde{\nabla}_{X_u}X_u &= X_{uu} = \Gamma_{11}^{1}X_u + \Gamma_{11}^{2}X_v + h(X_u, X_u), \\ \widetilde{\nabla}_{X_u}X_v &= X_{uv} = \Gamma_{12}^{1}X_u + \Gamma_{12}^{2}X_v + h(X_u, X_v), \\ \widetilde{\nabla}_{X_v}X_v &= X_{vv} = \Gamma_{22}^{1}X_u + \Gamma_{22}^{2}X_v + h(X_v, X_v). \end{aligned} \tag{12}$$

Taking in mind (4), (9) and (12) we get

$$\begin{aligned} h(X_u, X_u) &= X_{uu} + \frac{FF_u}{W^2}X_u - \frac{F_u}{W^2}X_v, \\ h(X_u, X_v) &= X_{uv} + \frac{FG_u}{2W^2}X_u - \frac{G_u}{2W^2}X_v, \\ h(X_v, X_v) &= X_{vv} + \frac{GG_u}{2W^2}X_u - \frac{FG_u}{2W^2}X_v. \end{aligned} \tag{13}$$

Consequently, by the use of (3), (5), (8) and (11) with (13) the Gaussian curvature and mean curvature vector of M become

$$K = -\frac{1}{2W}\left(\frac{G_u}{W}\right)_u, \tag{14}$$

and

$$\overrightarrow{H} = \frac{1}{2W^2}\left(Eh(X_v, X_v) - 2Fh(X_u, X_v) + Gh(X_u, X_u)\right), \tag{15}$$

respectively.

In the sequel, we consider some special cases;

Case I. Suppose

$$x_3 = \cos\varphi(u),\ x_4 = \sin\varphi(u), \tag{16}$$

then the position vector of the knotted sphere M can be represented as

$$X(u,v) = (x_1, x_2, \cos\varphi(u)\cos v - \sin\varphi(u)\sin v, \cos\varphi(u)\sin v + \sin\varphi(u)\cos v). \tag{17}$$

Hence, the coefficients of the first fundamental form become

$$\begin{aligned} E &= \langle X_u, X_u\rangle = 1, \\ F &= \langle X_u, X_v\rangle = \varphi'(u), \\ G &= \langle X_v, X_v\rangle = 1, \end{aligned} \tag{18}$$

where φ is a differentiable (angle) function.

Summing up (9)-(18) the following results are proved;

Proposition 3.1 *The surface M given with the position vector (17) is a flat surface.*

Proposition 3.2 *Let M be a surface given with the position vector (17). Then, the mean curvature $\left\|\overrightarrow{H}\right\|$ of M at point p is given by*

$$\left\|\overrightarrow{H}\right\| = \frac{1}{4\left(1-(\varphi'(u))^2\right)^2}\left(\kappa_\gamma^2 + 1 - 2(\varphi'(u))^2 - \frac{(\varphi''(u))^2}{1-(\varphi'(u))^2}\right), \tag{19}$$

where κ_γ is the curvature of the profile curve γ.

Proof 3.3 *With the help of (13), (15) and (18) the Gaussian curvature vector of M becomes*

$$2\overrightarrow{H} = \frac{1}{1-(\varphi')^2}\left(\overline{x}_1, \overline{x}_2, \overline{x}_3\cos v + \overline{x}_4\sin v, \overline{x}_3\sin v - \overline{x}_4\cos v\right), \tag{20}$$

where,

$$\begin{aligned}
\overline{x}_1 &= x_1'' + \frac{\varphi'\varphi''}{1-(\varphi')^2}x_1', \\
\overline{x}_2 &= x_1'' + \frac{\varphi'\varphi''}{1-(\varphi')^2}x_2' \\
\overline{x}_3 &= \frac{\varphi'\varphi''x_3' + \varphi''x_4}{1-(\varphi')^2} + x_3'' - x_3 + 2\varphi'x_4' = -\left(1-(\varphi')^2\right)\cos\varphi, \\
\overline{x}_4 &= \frac{\varphi''x_3 - \varphi'\varphi''x_4'}{1-(\varphi')^2} - x_4'' + x_4 + 2\varphi'x_3' = \left(1-(\varphi')^2\right)\sin\varphi
\end{aligned} \tag{21}$$

are differentiable functions. Taking the norm of the vector (20) and using (16) with (21) we obtain (19). This completes the proof of the proposition.

As a consequence of Proposition 2 we obtain the following result.

Corollary 3.4 *Let M be a surface given with the position vector (17). Then M is a minimal surface if and only if the curvature κ_γ of the profile curve γ satisfies the equality*

$$\kappa_\gamma^2 = \frac{(\varphi''(u))^2}{1-(\varphi'(u))^2} + 2\left(\varphi'(u)\right)^2 - 1, \tag{22}$$

in such a way that the (angle) function φ is non-constant.

Case II. Suppose $x_4 = \lambda x_3$, $\lambda \in \mathbb{R}$, then the position vector of the knotted sphere M can be represented as

$$r(s,\theta) = \left(x_1(u), x_2(u), x_3(u)\left(\cos v - \lambda \sin v\right), x_3(u)\left(\sin v + \lambda \cos v\right)\right). \tag{23}$$

Hence, the coefficients of the first fundamental form of M become

$$\begin{aligned}
E &= \langle X_u, X_u\rangle = 1, \\
F &= \langle X_u, X_v\rangle = 0, \\
G &= \langle X_v, X_v\rangle = \left(1+\lambda^2\right)x_3^2(u).
\end{aligned} \tag{24}$$

By the use of (24) with (3) we obtain the following result.

Proposition 3.5 *Let M be a surface given with the position vector (23). Then, the Gaussian curvature of M is given by*

$$K = -\frac{x_3''(u)}{x_3(u)}.$$

As a consequence of Proposition 4 we obtain the following result.

Corollary 3.6 *Let M be a surface given with the position vector (23). Then we have the following statements*

i) If $x_3(u) = ae^{cu} + be^{-cu}$ then the corresponding surface is pseudo-spherical, i.e., it has negative Gaussian curvature $K = -\frac{1}{c^2}$,

ii) If $x_3(u) = a\cos cu + b\sin cu$ then the corresponding surface is spherical, i.e., it has negative Gaussian curvature $K = \frac{1}{c^2}$, where a, b and c are real constants.

iii) If $x_3(u) = au + b$ then the corresponding surface is flat.

4 The Conjugate Nets and Laplace Transforms of Knotted Sphere

In the present section, we will give some basic relations of the conjugate net of curves on a surface in $\mathbb{R}^n$. A net of curves on a surface M is called conjugate, if at every point the tangent directions of the curves of the net separate harmonically the asymptotic directions. Taking the net to be parametric net with parameters u and v, the classical notion of the conjugate net usually can be stated in [11] as follows:

Definition 4.1 *Let M be a smooth surface given with the position vector $X : U \subset \mathbb{E}^2 \to \mathbb{E}^n$, and $N_1, ..., N_{n-2}$ normal vector fields of M in $\mathbb{E}^n$. If $X_{uv} = \frac{\partial^2 X}{\partial u \partial v}$ satisfies*

$$\langle X_{uv}, N_\alpha \rangle = 0, 1 \leq \alpha \leq n-2, \tag{25}$$

then (u, v) is called conjugate coordinates of X and the net woven by coordinate curves is called the conjugate net. For convenience, we denote the conjugate net by (u, v). Here, $\langle , \rangle$ denotes the inner product on $\mathbb{E}^n$ [11].

Equation (25) is equivalent to the condition that X_{uv} lies in the subspace spanned by X_u and X_v; i.e.,

$$X_{uv} = \Gamma_{12}^1 X_u + \Gamma_{12}^2 X_v. \tag{26}$$

Now, for the surface with normal conjugate net, we have two transforms

$$X_1 = X - \frac{X_v}{\Gamma_{12}^1},\ X_{-1} = X - \frac{X_u}{\Gamma_{12}^2}, \tag{27}$$

which are called the *Laplace transforms* of surface M [10].

Furthermore, the functions

$$h = \partial_u \Gamma_{12}^1 - \Gamma_{12}^1 \Gamma_{12}^2,\ k = \partial_v \Gamma_{12}^2 - \Gamma_{12}^1 \Gamma_{12}^2 \tag{28}$$

are called the *Laplace invariants*.

If $\Gamma_{12}^1 \neq 0$ (*resp.* $\Gamma_{12}^2 \neq 0$), the conjugate net is called $v-$*direction normal* (resp. $u-$*direction normal*). To establish geometrically the notion of conjugate net in ambient space, the following result explain the real geometric meaning of the conjugate net defined in (25).

Proposition 4.2 *[11] (u, v) is a v-direction normal conjugate net of M if and only if there exists another surface $\widetilde{M}$ given with the position vector $X_1(u, v)$ such that, for any $(u, v) \in D \subset \mathbb{E}^2$, the straight line XX_1 joining the points $X(u, v)$ and $X_1(u, v)$ is parallel to the vectors $X_v(u, v)$ and $X_{1u}(u, v)$.*

We obtain the following results,

Theorem 4.3 *Let M be a surface given with the position vector (6). If (u, v) are conjugate coordinates, then M is a flat surface.*

Proof 4.4 *Let (u, v) be conjugate coordinates of the knotted sphere M given with the parametrization (6). Then, by definition $h(X_u, X_v) = 0$. So, by the use of (13) we have*

$$X_{uv} = \frac{G_u}{2W^2} X_v - \frac{FG_u}{2W^2} X_u. \tag{29}$$

Consequently, substituting (7) with (10) into (29) we get

$$\begin{aligned} x_1'(u) &= 0, \\ x_2'(u) &= 0, \\ x_3'(u) &= x_3(u)\frac{G_u}{2W^2} - x_4'(u)\frac{FG_u}{2W^2} \\ x_4'(u) &= x_4(u)\frac{G_u}{2W^2} + x_3'(u)\frac{FG_u}{2W^2}. \end{aligned} \tag{30}$$

Summing up the last two equations of (30) we obtain

$$\left(x_3'(u)\right)^2 + \left(x_4'(u)\right)^2 = \frac{G_u}{2W^2}\left(x_3(u)x_3'(u) + x_4(u)x_4'(u)\right). \tag{31}$$

Moreover, the profile curve γ has arclength parameter and the first two equations of (30) imply that

$$\left(x_3'(u)\right)^2 + \left(x_4'(u)\right)^2 = 1. \tag{32}$$

Hence, by the use of (8) with (32) the equation (31) reduces to

$$1 = \frac{G_u^2}{4W^2},\ W > 0. \tag{33}$$

Thus, substituting (33) into (14) we get $K = -\frac{1}{2W}\left(\frac{G_u}{W}\right)_u = 0$. This completes the proof of the proposition.

By the virtue of (14) the following results hold.

Corollary 4.5 *The coordinates (u, v) of the surface M given with the position vector (17) can not be conjugate.*

Proof 4.6 *Suppose that (u, v) are the conjugate coordinates of the surface given with the parametrization (17). Then, from (18) and (33) we get $4W^2 = G_u^2 = 0$. But, this contradicts the fact that $W > 0$. So, the coordinates (u, v) can not be conjugate.*

Remark 4.7 *Corollary 7 shows that the inverse statement of Theorem 7 may not be true.*

Proposition 4.8 *Let M be a knotted sphere given with the position vector (23). Then the conjugate surface $\widetilde{M}$ is a part of the rotation plane Π.*

Proof 4.9 *Let M be a knotted sphere given with the parametrization (6), then by the use of (9) we get*

$$\Gamma_{12}^1 = -\frac{FG_u}{2W^2}, \Gamma_{12}^2 = \frac{G_u}{2W^2}.$$

Now, assume that the surface M with normal conjugate net, then equation (24) yields $\Gamma^1_{12} = 0$ and $\Gamma^2_{12} = \frac{x_3'(s)}{x_3(s)}$. Consequently, the Laplace transform X_{-1} becomes

$$X_{-1} = X - \frac{x_3(u)}{x_3'(u)} X_u.$$

Hence, using (23) with its partial derivative X_u we obtain

$$X_{-1} = \left(x_1(u) - \frac{x_3(u)}{x_3'(u)} x_1'(u), x_2(u) - \frac{x_3(u)}{x_3'(u)} x_2'(u), 0, 0 \right). \tag{34}$$

This completes the proof of the proposition.

References

[1] *Yu. A. Aminov*, Geometry of Submanifolds. Gordon & Breach Science Publ., Amsterdam, 2001.

[2] *E. Artin,* Zur izotopi zweidimensionaler Flachen im R4. *Abh. Math Sem. Univ. Hamburg*, **4**(1925), 174 − 177.

[3] *K. Arslan, B. Bayram, B. Bulca and G. Öztürk,* General rotation surfaces in E^4, *Results. Math.*, **61**(2012), 315 − 327.

[4] *B. Bulca, K. Arslan, B.K. Bayram and G. Öztürk,* Spherical product surfaces in E^4. *An. St. Univ. Ovidius Constanta*, **20**(2012), 41 − 54.

[5] *B.Y., Chen,* Geometry of Submanifolds, Dekker, New York, 1973.

[6] *D.V. Cuong,* Surfaces of Revolution with Constant Gaussian Curvature in Four- Space, arXiv:1205.2143v3.

[7] *U. Dursun and N.C. Turgay,* General rotational surfaces in Euclidean space E^4 with pointwise 1-type Gauss map . *Math. Commun.*, **17**(2012), 71 − 81.

[8] *G. Ganchev and V. Milousheva,* On the Theory of Surfaces in the Four-dimensional Euclidean Space. *Kodai Math. J.* **31**(2008), 183 − 198.

[9] *Gray, A.* Modern Differential Geometry of Curves and Surfaces. CRC Press, Boca Raton Ann Arbor London Tokyo, 1993.

[10] *N. Kamran and K. Tenenblat,* Laplace transformation in higher dimensions, *Duke Math. J.* **80**(1996), 237 − 266.

[11] *H. Li and Z. Guo,* The Conjugate Nets, Cartan Submanifolds, and Laplace Transformations in Space Forms1*, J. Math. Anal. Appl.* **267**(2002), 726 − 745.

[12] *C. Moore,* Surfaces of Rotation in a Space of Four Dimension. *Ann. Math.*, **21**(1919); 81 − 93.

[13] *Y.C. Wong,* Contributions to the theory of surfaces in 4-space of constant curvature, *Trans. Amer. Math. Soc.,* **59**(1946), 467 − 507.

Kadri Arslan
Department of Mathematics
Uludağ University
16059 Bursa, TURKEY
E-mail: arslan@uludag.edu.tr